

## Section 2.5

### LU Factorization

## First factorization of MATH 2802

A guru provides, for (suitable)  $m \times n$  matrix  $A$ , matrices  $L$  and  $U$  such that

- ▶  $L$  is **lower triangular**  $m \times m$  matrix (with *ones on the diagonal*)
- ▶  $U$  is an  $m \times n$  **row echelon form** (not necessary reduced)
- ▶  $A = LU$

E.g.

$$A = \begin{pmatrix} 2 & 4 & -1 & 5 & -1 \\ -4 & -5 & 3 & -8 & 1 \\ 2 & -5 & -4 & 1 & 8 \\ -6 & 0 & 7 & -3 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & -3 & 1 & 0 \\ -3 & 4 & 2 & 1 \end{pmatrix} \begin{pmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 5 \end{pmatrix}$$

The matrix  $LU$

It helps to interpret  $L$  as *instructions on how to sum rows* of  $U$ .

## A short cut to solving equations

**How to save time** in solving  $Ax = b$ ?

1. Visit the guru and *get  $L$  and  $U$* ,
2. *Quickly* solve  $Ly = b$ ,
3. *Quickly* solve  $Ux = y$ ,
4. Claim that  **$x$  is a solution** to  $Ax = b$ .

Are we **allowed to** do that?

Dense matrices -few entries are zeros-

Sparse matrices -most entries are zeros.

*Save up* to 2/3 of the time

If  $n \geq 30$ ,  $A$  is  $n \times n$  and dense

- ▶ Computing  $L$  and  $U$ :  $\sim 2n^3/3$  flops.
  - ▶ Solving a triangular system:  $\sim n^2$  flops
- vs.**
- ▶ Computing  $A^{-1}$ :  $\sim 2n^3$  flops
  - ▶ Matrix multiplication  $A^{-1}x$ :  $\sim 2n^2$
  - ▶ *Extra downside*: roundoff errors in  $A^{-1}$

Avoid dense matrices

If  $A$  is sparse it is likely that

- ▶  $A^{-1}$  is dense
- ▶  $L$  and  $U$  are sparse

## Become the guru and sell factorizations

**Our assumption:** Row reduction of  $A$  requires *no row swaps*

1. *Row reduce* the matrix  $A$ , but *do not normalize pivots* to being one's; e.g.

$$(E_6 E_5 E_4 E_3 E_2 E_1) A = U$$

2. Separate the row reduction according to 'clearing' pivot columns

$$(E_6)(E_5 E_4)(E_3 E_2 E_1) A = U$$

3. Now left multiply by the inverses of the elementary matrices:

$$A = ((E_6)(E_5 E_4)(E_3 E_2 E_1))^{-1} U = LU$$

Looks like a very complicated matrix:

$$L = ((E_6)(E_5 E_4)(E_3 E_2 E_1))^{-1}$$

but can be **read off the row-reduction steps**.

## Construct the matrix $L$

$$L = ((E_6)(E_5 E_4)(E_3 E_2 E_1))^{-1}$$

2. Separate the row reduction according to 'clearing' pivot columns

$$A = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ -4 & -5 & 3 & -8 & 1 \\ 2 & -5 & -4 & 1 & 8 \\ -6 & 0 & 7 & -3 & 1 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & -9 & -3 & -4 & 10 \\ 0 & 12 & 4 & 12 & -5 \end{bmatrix} = A_1$$

$$\sim A_2 = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 4 & 7 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 5 \end{bmatrix} = U$$

$$\begin{bmatrix} 2 \\ -4 \\ 2 \\ -6 \end{bmatrix} \begin{bmatrix} 3 \\ -9 \\ 12 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \end{bmatrix} \begin{bmatrix} 5 \end{bmatrix}$$

$$\div 2 \quad \div 3 \quad \div 2 \quad \div 5$$



$$\begin{bmatrix} 1 & & & \\ -2 & 1 & & \\ 1 & -3 & 1 & \\ -3 & 4 & 2 & 1 \end{bmatrix},$$

$$\text{and } L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & -3 & 1 & 0 \\ -3 & 4 & 2 & 1 \end{bmatrix}$$

## Construct the matrix $L$

$$L = ((E_6)(E_5 E_4)(E_3 E_2 E_1))^{-1}$$

2. Separate the row reduction according to 'clearing' pivot columns

$$A = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ -4 & -5 & 3 & -8 & 1 \\ 2 & -5 & -4 & 1 & 8 \\ -6 & 0 & 7 & -3 & 1 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & -9 & -3 & -4 & 10 \\ 0 & 12 & 4 & 12 & -5 \end{bmatrix} = A_1$$

$$\sim A_2 = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 4 & 7 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 5 \end{bmatrix} = U$$

$$\begin{bmatrix} 2 \\ -4 \\ 2 \\ -6 \end{bmatrix} \begin{bmatrix} 3 \\ -9 \\ 12 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \end{bmatrix} \begin{bmatrix} 5 \end{bmatrix}$$

$$\div 2 \quad \div 3 \quad \div 2 \quad \div 5$$



$$\begin{bmatrix} 1 & & & \\ -2 & 1 & & \\ 1 & -3 & 1 & \\ -3 & 4 & 2 & 1 \end{bmatrix},$$

$$\text{and } L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & -3 & 1 & 0 \\ -3 & 4 & 2 & 1 \end{bmatrix}$$

## Process on the backstage

This backstage process can be *read off the row-reduction steps*.

1. *Divide into streaks* of elementary matrices with non-zero entries on the same column. E.g.  $E_3E_2E_1$
2. *Superposition*: The product of these elementary matrices will have the same non-entries (see atomic triangular matrix).
3. *Inverse*:  $(E_3E_2E_1)^{-1}$  changes the sign of all off-diagonal entries.

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 3 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 3 & 0 & 0 & 1 \end{pmatrix}$$
$$(E_3E_2E_1)^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ -3 & 0 & 0 & 1 \end{pmatrix}$$



## A second example

Find the  $LU$  factorization of  $A$ :

$$A = \begin{bmatrix} 2 & -4 & -2 & 3 \\ 6 & -9 & -5 & 8 \\ 2 & -7 & -3 & 9 \\ 4 & -2 & -2 & -1 \\ -6 & 3 & 3 & 4 \end{bmatrix} \sim \begin{bmatrix} 2 & -4 & -2 & 3 \\ 0 & 3 & 1 & -1 \\ 0 & -3 & -1 & 6 \\ 0 & 6 & 2 & -7 \\ 0 & -9 & -3 & 13 \end{bmatrix} \rightarrow \begin{bmatrix} 2 \\ 6 \\ 2 \\ 4 \\ -6 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \\ 6 \\ -9 \end{bmatrix} \begin{bmatrix} 5 \\ -5 \\ 10 \end{bmatrix}$$

$\div 2 \quad \downarrow \quad \div 3 \quad \downarrow \quad \div 5 \quad \downarrow$

$$\sim \begin{bmatrix} 2 & -4 & -2 & 3 \\ 0 & 3 & 1 & -1 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & -5 \\ 0 & 0 & 0 & 10 \end{bmatrix} \sim \begin{bmatrix} 2 & -4 & -2 & 3 \\ 0 & 3 & 1 & -1 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$



$$\begin{bmatrix} 1 & & & & \\ 3 & 1 & & & \\ 1 & -1 & 1 & & \\ 2 & 2 & -1 & 1 & \\ -3 & -3 & 2 & 0 & 1 \end{bmatrix},$$

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 & 0 \\ 1 & -1 & 1 & 0 & 0 \\ 2 & 2 & -1 & 1 & 0 \\ -3 & -3 & 2 & 0 & 1 \end{bmatrix}$$

# Electrical networks

Electric networks can be analysed as composition of smaller networks.

$$\begin{pmatrix} v_2 \\ i_2 \end{pmatrix} = A \begin{pmatrix} v_1 \\ i_1 \end{pmatrix}$$



**Transfer matrix  $A$ :** relation between the *input*  $\begin{pmatrix} v_2 \\ i_2 \end{pmatrix}$  and the *output*  $\begin{pmatrix} v_1 \\ i_1 \end{pmatrix}$ .

Engineers gives the following information:

$$\begin{bmatrix} 1 & -R_1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 0 \\ -1/R_2 & 1 \end{bmatrix}$$

Transfer matrix of series circuit      Transfer matrix of shunt circuit

## A ladder circuit

If  $\begin{pmatrix} v_1 \\ i_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  and the resistances in the circuit are given by  $R_1 = 3$  and  $R_2 = .5$ . Give the value of  $\begin{pmatrix} v_3 \\ i_3 \end{pmatrix}$

