

Announcements

Monday, October 23

- ▶ **Webwork** due next week,
- ▶ No quiz this week.

Chapter 3

Determinants

Section 3.1

Introduction to Determinants

Orientation

Recall: This course is about learning to:

- ▶ Solve the matrix equation $Ax = b$
We've said most of what we'll say about this topic now.
- ▶ Solve the matrix equation $Ax = \lambda x$ (eigenvalue problem)
We are now aiming at this.
- ▶ Almost solve the equation $Ax = b$
This will happen later.

The next topic is **determinants**.

This is a completely *magical function* that takes a square matrix and gives you a number.


It is a very complicated functionthe formula for the determinant of a 10×10 matrix has 3,628,800 summands! so we need efficient ways to compute it.

Today is mostly about the *computation* of determinants; in the next lecture we will focus on the *theory*.

The Idea of Determinants

Let A be an $n \times n$ matrix. **Determinants are only for square matrices.**

The columns v_1, v_2, \dots, v_n give you n vectors in \mathbf{R}^n . These determine a **parallelepiped** P .



Observation: the volume of P is zero \iff the columns are *linearly dependent* (P is “flat”) \iff the matrix A is not invertible.

The **determinant** of A will be a number $\det(A)$ whose absolute value is the *volume of P* . In particular, $\det(A) \neq 0 \iff A$ is invertible.


Determinants of 2×2 Matrices

Revisited

We already have a formula in the 2×2 case:

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc.$$

What does this have to do with volumes?



$$v_1 = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

The area of the parallelogram is

$$\text{base} \times \text{height} = 2 \cdot 3 = \left| \det \begin{pmatrix} 2 & 1 \\ 0 & 3 \end{pmatrix} \right|.$$

The area of the parallelogram is always $|ad - bc|$. If v_1 is not on the x -axis: it's a fun geometry problem! **Note:** this shows $\det(A) \neq 0 \iff A$ is invertible in this case. (The volume is zero if and only if the columns are collinear.)
Question: What does the sign of the determinant mean?

Determinants of 3×3 Matrices

Here's the formula:

$$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{aligned} & a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ & - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} \end{aligned}$$

How to remember this?

Draw a bigger matrix, repeating the first two columns to the right:

$$+ \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{vmatrix} - \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{vmatrix}$$

Then add the products of the downward diagonals, and subtract the product of the upward diagonals.

For example,

$$\det \begin{pmatrix} 5 & 1 & 0 \\ -1 & 3 & 2 \\ 4 & 0 & -1 \end{pmatrix} = \begin{vmatrix} 5 & 1 & 0 & 5 & 1 \\ -1 & 3 & 2 & -1 & 3 \\ 4 & 0 & -1 & 4 & 0 \end{vmatrix} = -15 + 8 + 0 - 0 - 0 - 1 = -8$$

What does this have to do with *volumes*? *Next time.*

A Formula for the Determinant

When $n \geq 4$, the determinant is **not that simple to describe**. The formula is recursive: you compute a larger determinant in terms of smaller ones.

We need some notation. Let A be an $n \times n$ matrix.

A_{ij} = ij th **minor** of A

= $(n-1) \times (n-1)$ matrix you get by *deleting the i th row and j th column*

C_{ij} = ij th **cofactor** of $A = (-1)^{i+j} \det A_{ij}$

The signs of the cofactors follow a *checkerboard pattern*:

$$\begin{pmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{pmatrix} \quad \pm \text{ in the } ij \text{ entry is the sign of } C_{ij}$$

Definition

The **determinant** of an $n \times n$ matrix A is

$$\det(A) = \sum_{j=1}^n a_{1j} C_{1j} = a_{11} C_{11} + a_{12} C_{12} + \cdots + a_{1n} C_{1n}.$$

This formula is called **cofactor expansion** *along the first row*.

A Formula for the Determinant

1×1 Matrices

This is the beginning of the recursion.

$$\det(a_{11}) = a_{11}.$$

A Formula for the Determinant

2×2 Matrices

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

The minors are:

$$A_{11} = \begin{pmatrix} \cancel{a_{11}} & \cancel{a_{12}} \\ \cancel{a_{21}} & a_{22} \end{pmatrix} = (a_{22})$$

$$A_{12} = \begin{pmatrix} \cancel{a_{11}} & \cancel{a_{12}} \\ a_{21} & \cancel{a_{22}} \end{pmatrix} = (a_{21})$$

$$A_{21} = \begin{pmatrix} \cancel{a_{11}} & a_{12} \\ \cancel{a_{21}} & \cancel{a_{22}} \end{pmatrix} = (a_{12})$$

$$A_{22} = \begin{pmatrix} a_{11} & \cancel{a_{12}} \\ \cancel{a_{21}} & \cancel{a_{22}} \end{pmatrix} = (a_{11})$$

The cofactors are

$$C_{11} = + \det A_{11} = a_{22}$$

$$C_{12} = - \det A_{12} = -a_{21}$$

$$C_{21} = - \det A_{21} = -a_{12}$$

$$C_{22} = + \det A_{22} = a_{11}$$

The determinant is

$$\det A = a_{11}C_{11} + a_{12}C_{12} = a_{11}a_{22} - a_{12}a_{21}.$$

A Formula for the Determinant

3 × 3 Matrices

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

The top row minors and cofactors are:

$$A_{11} = \begin{pmatrix} \cancel{a_{11}} & \cancel{a_{12}} & \cancel{a_{13}} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix} \quad C_{11} = + \det \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix}$$

$$A_{12} = \begin{pmatrix} \cancel{a_{11}} & \cancel{a_{12}} & \cancel{a_{13}} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{pmatrix} \quad C_{12} = - \det \begin{pmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{pmatrix}$$

$$A_{13} = \begin{pmatrix} \cancel{a_{11}} & \cancel{a_{12}} & \cancel{a_{13}} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix} \quad C_{13} = + \det \begin{pmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\det A = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$

$$= a_{11} \det \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix} - a_{12} \det \begin{pmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{pmatrix} + a_{13} \det \begin{pmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

The determinant is the same formula as before (*verify it yourself*)

A Formula for the Determinant

Example

$$\begin{aligned}\det \begin{pmatrix} 5 & 1 & 0 \\ -1 & 3 & 2 \\ 4 & 0 & -1 \end{pmatrix} &= 5 \cdot \det \begin{pmatrix} \cancel{5} & \cancel{1} & \cancel{0} \\ -\cancel{1} & -3 & 2 \\ \cancel{4} & 0 & -1 \end{pmatrix} - 1 \cdot \det \begin{pmatrix} \cancel{5} & \cancel{1} & \cancel{0} \\ -1 & \cancel{3} & 2 \\ 4 & \cancel{0} & -1 \end{pmatrix} \\ &\quad + 0 \cdot \det \begin{pmatrix} \cancel{5} & \cancel{1} & \cancel{0} \\ -1 & 3 & \cancel{2} \\ 4 & 0 & \cancel{-1} \end{pmatrix} \\ &= 5 \cdot \det \begin{pmatrix} 3 & 2 \\ 0 & -1 \end{pmatrix} - 1 \cdot \det \begin{pmatrix} -1 & 2 \\ 4 & -1 \end{pmatrix} + 0 \cdot \det \begin{pmatrix} -1 & 3 \\ 4 & 0 \end{pmatrix} \\ &= 5 \cdot (-3 - 0) - 1 \cdot (1 - 8) \\ &= -15 + 7 = -8\end{aligned}$$

Cofactor expansion: Specify point of reference...

Recall: the formula

$$\det(A) = \sum_{j=1}^n a_{1j} C_{1j} = a_{11} C_{11} + a_{12} C_{12} + \cdots + a_{1n} C_{1n}.$$

is called cofactor expansion *along the first row*. Actually, you can expand cofactors along any **row** or **column** you like!

$$\det A = \sum_{j=1}^n a_{ij} C_{ij} \quad \text{for any fixed } i$$

$$\det A = \sum_{i=1}^n a_{ij} C_{ij} \quad \text{for any fixed } j$$

Good trick: Use cofactor expansion along a row or a column *with a lot of zeros*.

Cofactor Expansion

Example

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 1 & 0 \\ 5 & 9 & 1 \end{pmatrix}$$

It looks easiest to expand along the third column:

$$\begin{aligned} \det A &= 0 \cdot \det \begin{pmatrix} \text{don't} \\ \text{care} \end{pmatrix} - 0 \cdot \det \begin{pmatrix} \text{don't} \\ \text{care} \end{pmatrix} + 1 \cdot \det \begin{pmatrix} 2 & 1 & 0 \\ 1 & 1 & 0 \\ 5 & 9 & 1 \end{pmatrix} \\ &= \det \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} = 2 - 1 = 1 \end{aligned}$$

Poll

$$\det \begin{pmatrix} 1 & 7 & -5 & 14 & 3 & 22 \\ 0 & -2 & -3 & 13 & 11 & 1 \\ 0 & 0 & -1 & -9 & 7 & 18 \\ 0 & 0 & 0 & 3 & 6 & -8 \\ 0 & 0 & 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix} = ?$$

A. -6 B. -3 C. -2 D. -1 E. 1 F. 2 G. 3 H. 6

If you *expand repeatedly along the first column*, you get

$$\begin{aligned} 1 \cdot \det \begin{pmatrix} -2 & -3 & 13 & 11 & 1 \\ 0 & -1 & -9 & 7 & 18 \\ 0 & 0 & 3 & 6 & -8 \\ 0 & 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix} &= 1 \cdot (-2) \cdot \det \begin{pmatrix} -1 & -9 & 7 & -18 \\ 0 & 3 & 6 & -8 \\ 0 & 0 & 1 & -11 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ &= 1 \cdot (-2) \cdot (-1) \cdot \det \begin{pmatrix} 3 & 6 & -8 \\ 0 & 1 & -11 \\ 0 & 0 & -1 \end{pmatrix} = 1 \cdot (-2) \cdot (-1) \cdot 3 \cdot \det \begin{pmatrix} 1 & -11 \\ 0 & -1 \end{pmatrix} \\ &= 1 \cdot (-2) \cdot (-1) \cdot 3 \cdot 1 \cdot (-1) = -6 \end{aligned}$$

The Determinant of an Upper-Triangular Matrix

Trick: Expand along the first row

This works for any matrix that is *upper-triangular* (all entries below the main diagonal are zero).

Theorem

The determinant of an upper-triangular matrix is the product of the diagonal entries:

$$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_{nn} \end{pmatrix} = a_{11} a_{22} a_{33} \cdots a_{nn}.$$

The same is true for lower-triangular matrices. (Repeatedly expand along the first row.)

Extra: A Formula for the Inverse


from §3.3

For 2×2 matrices we had a nice formula for the inverse:

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \implies A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \frac{1}{\det A} \begin{pmatrix} C_{11} & C_{21} \\ C_{12} & C_{22} \end{pmatrix}.$$

Theorem

This last formula works for any $n \times n$ invertible matrix A :


$$A^{-1} = \frac{1}{\det A} \begin{pmatrix} C_{11} & C_{21} & C_{31} & \cdots & C_{n1} \\ C_{12} & C_{22} & C_{32} & \cdots & C_{n2} \\ C_{13} & C_{23} & C_{33} & \cdots & C_{n3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ C_{1n} & C_{2n} & C_{3n} & \cdots & C_{nn} \end{pmatrix} = \frac{1}{\det A} (C_{ij})^T$$

Note that the cofactors are “transposed”: the (i, j) entry of the matrix is C_{ji} .

The proof uses Cramer’s rule. See Dan Margalit’s notes (p. 64) on the website for a nice explanation.

A Formula for the Inverse

Example

Compute A^{-1} , where $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}$.

The minors are:

$$A_{11} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad A_{12} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad A_{13} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

$$A_{21} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad A_{22} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad A_{23} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$

$$A_{31} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \quad A_{32} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad A_{33} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

The cofactors are (don't forget to multiply by $(-1)^{i+j}$):

$$C_{11} = -1 \quad C_{12} = 1 \quad C_{13} = -1$$

$$C_{21} = 1 \quad C_{22} = -1 \quad C_{23} = -1$$

$$C_{31} = -1 \quad C_{32} = -1 \quad C_{33} = 1$$

The determinant is (expanding along the first row):

$$\det A = 1 \cdot C_{11} + 0 \cdot C_{12} + 1 \cdot C_{13} = -2$$

Extra: A Formula for the Inverse

Example, continued

Compute A^{-1} , where $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}$.

The inverse is

$$A^{-1} = \frac{1}{\det A} \begin{pmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} -1 & 1 & -1 \\ 1 & -1 & -1 \\ -1 & -1 & 1 \end{pmatrix}.$$

Check:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix} \cdot -\frac{1}{2} \begin{pmatrix} -1 & 1 & -1 \\ 1 & -1 & -1 \\ -1 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad \checkmark$$